

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Triclopyr 600

HERBICIDE

ACTIVE CONSTITUENT: 600 g/L TRICLOPYR present as the BUTOXYETHYL ESTER

GROUP I HERBICIDE

For the control of a range of woody weeds and melons as specified in the Directions for Use table.

CONTENTS

20 Litres

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE. SHAKE WELL BEFORE USE.

Australis Crop Protection Pty Ltd

ABN 94 150 711 185

Shop 4, 30 Heber Street, Moree NSW 2400

Telephone: 0417 329 133 • Facsimile: 07 3337 9882

www.austcrop.com.au

BATCH NO:

DATE OF MANUFACTURE:



Triclopyr 600

HERBICIDE

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area as cool as possible. DO NOT store for prolonged periods in direct sunlight. DO NOT store with seeds, foodstuffs or fertilizer. DO NOT use this container for any other purpose.

For Non-Refillable Containers: Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Australis Crop Protection Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid as above for disposal.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length rubber gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Triclopyr 600 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval No.: 66566/53900

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE



AUSTRALIS
CROP PROTECTION

Australis Crop Protection Pty Ltd
ABN 94 150 711 185

Shop 4, 30 Heber Street, Moree NSW 2400
Telephone: 0417 329 133 • Facsimile: 07 3337 9882

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Triclopyr 600

HERBICIDE

ACTIVE CONSTITUENT: 600 g/L TRICLOPYR
present as the BUTOXYETHYL ESTER

GROUP I HERBICIDE

For the control of a range of woody weeds and melons as
specified in the Directions for Use table.

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No.: 65548/51227



Australis Crop Protection Pty Ltd
ABN 94 150 711 185
Shop 4, 30 Heber Street, Moree NSW 2400
Telephone: 0417 329 133 • Facsimile: 07 3337 9882
www.austcrop.com.au

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area as cool as possible. DO NOT store for prolonged periods in direct sunlight. DO NOT store with seeds, foodstuffs or fertilizer. DO NOT use this container for any other purpose.

For Non-Refillable Containers: Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Australis Crop Protection Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid as above for disposal.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length rubber gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Triclopyr 600 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.



PEEL HERE

DIRECTIONS FOR USE:

Restrains: DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.

DO NOT spray if rain is likely within one (1) hour or if foliage is wet from rain or dew. However, when tankmixed with Glyphosate 450 g/L, this time extends to six (6) hours.

DO NOT burn off, cut or clear blackberry or other woody weeds for six (6) months after spraying.

1. WOODY WEEDS SITUATIONS**Table A: High volume spraying**

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100L WATER	CRITICAL COMMENTS
African Boxthorn	Less than 2m tall	Tas only	170 mL	
Angophora sp. Banksia sp.	1 to 2m tall	All States	400 or 560 mL	Use the higher rate on larger regrowth. Ensure the weed has dense foliage.
Blackberry in association with: St John's Wort	Active growth during late spring to early autumn		170 mL	Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before spraying with this product. Any subsequent regrowth and seedlings should be sprayed after hardening off.
	During flowering (Nov-Jan)	NSW, Vic, Tas only		Apply as a thorough foliage spray.
Brigalow	1 to 2m tall	Qld, NSW only		Use at least 1000 L of water/ha.
Brooms: English	Spring to mid-summer prior to pod formation	All States		
Camphor Laurel	Seedlings up to 3m tall			
Capeweed	Rosette	Tas only	80 mL	
Common Prickly Pear	Active growth	All States	3 L	
English Ivy	Active growth during late spring to late summer	Vic only	1 L + 1 L glyphosate (360 g/L)	DO NOT treat ivy growing up trees or on other plants as death of the host may result. This mixture is not selective to grasses.
Eucalyptus	Seedlings and regrowth for small lignotubers, 1 to 2m tall	Qld, SA, WA, NT only	400 mL	Add Decol [®] S35B Surfactant at 250 mL to 500 mL/100 L of water for best results.
		NSW, Vic, Tas only	560 mL	
Fennel	1 to 2m tall	Tas only	170 mL	Some regrowth may be expected the following season which can be sprayed after hardening off.
Green Cestrum		Qld, NSW, Vic only		
Groundsel Bush (<i>Baccharis halimifolia</i>)	Seedlings, 1 to 2m tall	All States	160 mL	
	2 to 3m tall		320 mL	
Gorse	Spring to mid summer		170 mL or 340 mL	Add a 100% concentrate non-ionic wetting agent at rate of 125 mL/100L water. Retreatment of regrowth may be necessary. Use higher water rate on older hardened off plants.

Table A: High volume spraying – continued

WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/100L WATER	CRITICAL COMMENTS
Horehound	Rosette	Tas only	170 mL	
Saffron Thistle	Up to bud stage		80 mL	
Tiger Pear	Active growth	All States	3 L	
Wattles including Silver Wattle	Seedlings, 1 to 2m tall		160 mL	
Black Wattle	2 to 3m tall		320 mL	

Table B: Aerial application

Helicopter NSW, SA, Tas, Vic and WA only

Helicopter or fixed wing aircraft (Qld only)

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / Ha	CRITICAL COMMENTS
Blackberry	Late spring to autumn	All States	4.8 L	AVOID overspray/drift onto waterways

Table C: Controlled Droplet Application (C.D.A.)

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/1L WATER	CRITICAL COMMENTS
Blackberry	Late spring to autumn	All States	170 mL	

Table D: Low Volume High Concentrate Application Techniques (Gas Powered Gun, Sprinkler Sprayer)

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/10L WATER	CRITICAL COMMENTS
Blackberry	Late spring to autumn	All States	280 mL	
Eucalypt seedlings	1 to 2m tall		400 mL	

Table E: Basal Bark and Cut Stump Treatment

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/60 L DIESEL	CRITICAL COMMENTS
African Boxthorn	Basal Bark: Plants up to 5cm basal diameter	All States	2 L	
Australian Blackthorn			1 L	
Broom		Tas only	1.25 L	
Brown Salwood	Cut Stump: Plants up to and in excess of basal bark sizes	All States	0.5 L	
Bitter bark		Qld, NSW only	1 L	
Castor Oil, Chinee Apple		All States		
Dawson gum		Qld only	2 L	
Eucalyptus (except Dawson Gum) False Sandalwood Green Wattle Lantana Needlewood		All States	1 L	
Olive		SA only	2 L	
		NSW only	4 L	
Paperbark Teatree Rubbervine Silver Wattle		All States	1 L	
Sweet Briar			2 L	
Yellow-wood		Qld only		
Camphor Laurel	Basal bark:	Qld, NSW only	1 L	
Common Prickly Pear	Plants up to 10cm basal diameter	All States	0.8 L	Apply as a thorough foliar spray.
Groundsel Bush (<i>Baccharis halimifolia</i>)	Cut stump: Plants up to and in excess of basal bark sizes		0.5 L	Treat from early summer rains to end of April when regrowth is apparent.
Prickly Acacia				
Privet			5 L	Treatment may be carried out at any time of the year.
Smooth Tree Pear Tiger Pear			0.8 L	Apply as a thorough foliage spray.
Tree of heaven			1 L	

Table E: Basal Bark and Cut Stump Treatment - continued

FENCELINES AND FIRE TRAILS ONLY				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE/60 L DIESEL	CRITICAL COMMENTS
Broadleaf Hopbush, Narrowleaf Hopbush, Turpentine Bush	Basal bark: Plants up to 10cm basal diameter	NSW only	1 L	

2. CROPPING/FALLOW SITUATIONS

See **GENERAL INSTRUCTIONS – APPLICATION** section for application method details.

FALLOW, STUBBLE, FIREBREAKS				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / HA	CRITICAL COMMENTS
Camel (Afghan, Bitter) Melon (<i>Citrullus lanatus</i>)	Up to 20cm diameter	Qld, NSW, Vic, SA, WA only	120 mL**	There are some strains of melon that are not controlled. Contact your Australis Crop Protection representative for more information. ** Add a crop oil such as Uptake [†] Spraying Oil at 500 mL/100 L water or D-C-Trate [†] Crop Oil at 1 L/100 L water. DO NOT use oils when tank mixing with Glyphosate herbicide. See COMPATIBILITY section. This mixture is not selective to grasses.
	Runners from 20 to 40 cm diameter		160 mL**	
Prickly Paddy Melon (<i>Cucumis myriocarpus</i>)	Up to 20cm diameter		80 mL**	
	Runners from 20 to 40cm diameter		160 mL**	

SORGHUM (Apply between 4 to 6 leaf stage, when secondary roots have developed)				
WEEDS CONTROLLED	WEED GROWTH STAGE	STATE	RATE / HA	CRITICAL COMMENTS
Prickly Paddy Melon	Up to 20cm diameter	Qld, NSW only	80 mL	DO NOT add crop oils, as severe crop damage may occur. Fusing of sorghum prop roots may be observed and this may be worse under stress (e.g. root disease). Should only be mixed with Starane [†] 200 Herbicide and atrazine (500 g/L flowable only) for increased weed spectrum.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA FOR BLACKBERRY:

DO NOT treat bushes carrying mature or near mature fruit.

FOR NATIVE VEGETATION:

Use of ACP Triclopyr 600 Herbicide on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

WITHHOLDING PERIODS:

PASTURE: NOT REQUIRED WHEN USED AS DIRECTED.

SORGHUM: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

COMPATIBILITY

Fallow Situations: ACP Triclopyr 600 Herbicide is compatible with the following products: Lontrel[†] Herbicide, Starane[†] 200 Herbicide, Glyphosate 450 g/L Herbicide, Glyphosate 490 g/L Herbicide, Touchdown[†] Broadacre, 2,4-D amine or 2,4-D ester, Imidacloprid 200 g/L Insecticide.

When mixing with Glyphosate 450 g/L in fallow, refer to the Glyphosate 450 label for use rate and adjuvants recommended.
DO NOT use Uptake[†] Spraying Oil or D-C-Trate Crop Oil.

Sorghum: ACP Triclopyr 600 Herbicide is compatible with Starane[†] 200 Herbicide and 2,4-D Amine 500 g/L Herbicide.

MINIMUM RECROPPING PERIODS

Before using this product in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the product with the longest plant-back period.

Observe the following recropping periods for ACP Triclopyr 600 Herbicide:

Wheat, barley, sorghum, maize	7 days
Chickpeas, soybeans, sunflowers	7 days
Cotton	14 days

MIXING

Half fill the spray unit with water and add the required amount of ACP Triclopyr 600 Herbicide. Add the remaining water with the agitator running. If required, then add crop oils or wetter (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

Basal Bark and Cut Stump Application: Quarter fill the spray unit or mixing container with diesel and add the required amount of ACP Triclopyr 600 Herbicide. Add the remaining diesel and shake or agitate thoroughly to mix the contents. Periodically shake or agitate to stop product settling out. Only mix sufficient solution for immediate daily use and avoid storing.

APPLICATION

1. WOODY WEED SITUATIONS

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to one metre high in situations which have been bulldozed, slashed, burnt, ploughed or areas have a previous chemical treatment.

A. High Volume Spraying: Thorough coverage of foliage and stems to the point of runoff is essential, however, avoid excess spraying which is wasteful of chemical.

Hand Gun: Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa. A spray volume of 3,000 to 4,000 L per infested hectare (30 to 40 L/100 m²) should be used on the weed infestation.

Knapsack: Used for smaller infestations, where penetration of the canopy is not essential. A spray volume of 3 to 4 L/10 m² of infested area should be used.

B. Aerial Application

Apply in 100 to 200 L water/ha. Use a calibrated aircraft to apply in half overlap passes. Nozzle configurations should produce droplets of not less than 250 to 350 micron diameter (e.g. D8/45). The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.

DO NOT spray when wind exceeds 10 km/hr and/or air temperature reaches 30°C.

C. Controlled Droplet Application (C.D.A.)

Results similar to high volume spraying can be obtained using Micron Herbi or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm². Use a marking agent as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A. unit.

D. Low Volume High Concentrate Application Techniques

Good control will be achieved, similar to high volume application, where bush size enables good coverage of the bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

Gas powered gun: Apply 50 mL shots to obtain uniform coverage of 4 to 5 m² of surface area of bush. This relates to 20 droplets/cm² of leaf surface.

Sprinkler sprayer: This technique involves using a micro sprinkler which is connected to a hollow fibreglass rod attached to a pressure knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

E. Basal Bark and Cut Stump Treatment

Mix ACP Triclopyr 600 Herbicide in diesel. The use of diesel as a herbicide carrier may affect the rubber seals in some sprayers. To avoid this, use sprayers which use Viton[†] seals and fittings.

Basal Bark method: DO NOT apply to wet stems as this can repel the diesel mixture.

Apply only with hand-directed equipment such as a pressure sprayer or a paint brush. Spray equipment should be used at low pressures, up to 200 kPa, to avoid excessive splashing or drift. Species with old, rough bark require more thorough wetting than smooth barked species. Liberally spray or paint the bark around the stem from ground level up to 30 cm high, wetting thoroughly to the point of runoff (unless otherwise stated).

Cut Stump Method: Stems should be cut less than 15 cm above the ground. **Immediately** apply ACP Triclopyr 600 Herbicide/diesel mixture **liberally** to the **freshly** cut stump by spray or painting the cut surface and sides of the stem.

2. CROPPING/FALLOW SITUATIONS

A. Boom Application

Application of ACP Triclopyr 600 Herbicide in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressures in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

B. Aerial Application

DO NOT allow ACP Triclopyr 600 Herbicide to physically drift onto desirable plants.

Aircraft may be used to apply ACP Triclopyr 600 Herbicide in fallow situations, when ground application equipment cannot be used due to prolonged wet conditions.

A minimum spray volume of 35 L/ha should be used with flat fan and CP nozzle configurations.

DO NOT apply ACP Triclopyr 600 by aircraft when wind exceeds 10 km/hr and/or air temperature reaches 30°C.

Droplets with an average size (or Volume Mean Diameter) of 250 to 350 micron diameter are recommended.

CLEANING SPRAY EQUIPMENT

Water Based Cleaning

Rinsing: After using ACP Triclopyr 600 Herbicide, empty the spray unit completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose.

Drain the spray unit and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination: Before spraying cotton and other sensitive crops with equipment that has been used to apply ACP Triclopyr 600 Herbicide, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section. Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF, OMO, DRIVE at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water) and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and water courses.

Diesel Based Cleaning

Rinsing: After using ACP Triclopyr 600 Herbicide, empty the spray unit completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose.

Drain the spray unit and clean any filters in the tank, pump, lines, hoses and nozzles. On completion of spraying, use a degreaser such as Caltex Kwik-D-Grease to remove traces of diesel from the sprayer. Rinse tank and spray through the nozzles with water to remove degreaser.

Decontamination: After the above, quarter fill the tank with clean water and add an alkali detergent (e.g. liquid SURF, OMO, OMOMATIC, DRIVE at 50 mL/10 L of water or the powder equivalent at 50 g/10 L of water). Shake or operate spray to circulate the washing solution throughout the sprayer and spray the solution through the nozzle. Rinse well with clean water to remove detergent. To clean brushes and container, spray liberally with degreaser. Hose off thoroughly with clean water and repeat using detergents (see above). DO NOT use this equipment for any other purpose.

RESISTANT WEEDS WARNING

ACP Triclopyr 600 Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I Herbicide. Some naturally occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other disrupters of plant cell growth herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Australis Crop Protection Pty Ltd accepts not liability for any losses that may result from the failure of this product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical suppliers, consultant, local Department of Agriculture, or Australis Crop Protection representative.

GROUP	I	HERBICIDE
-------	---	-----------

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to ACP Triclopyr 600 Herbicide include, but are not limited to: peas, lupins, Lucerne, navy beans, soybeans and other legumes; cotton, fruit, hops, ornamentals, shade trees and Pinus spp., potatoes, safflower, sugarbeet, sunflower, tobacco, tomatoes, vegetables and vines.

ACP Triclopyr 600 Herbicide is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment. DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops. When using ACP Triclopyr 600 Herbicide and Glyphosate 450 by aerial application in fallow situations, observance of a buffer zone of 150 metres to protect native tree species is required.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is highly toxic to fish and other aquatic organisms. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area as cool as possible. DO NOT store for prolonged periods in direct sunlight. DO NOT store with seeds, foodstuffs or fertilizer. DO NOT use this container for any other purpose.

For Non-Refillable Containers: Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs Australis Crop Protection Pty Ltd should be advised immediately. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid as above for disposal.

SAFETY DIRECTIONS

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin. When opening container and preparing spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow length rubber gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor, or Poisons Information Centre. Phone Australia 13 11 26.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE: The use of ACP Triclopyr 600 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Australis Crop Protection Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Australis Crop Protection Pty Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

† Other registered trademarks